See discussions, stats, and author profiles for this publication at: https://www.researchgate.net/publication/348980789

Theoretical investigation of X12O12 (X=Be, Mg, and Ca) in sensing CH2N2: A DFT study

Article in Computational and Theoretical Chemistry · February 2021

DOI: 10.1016/j.comptc.2021.113168

CITATIONS 0	5	READS 36	
4 authoi	rs, including:		
	Mohsen Doust Mohammadi 21 PUBLICATIONS 57 CITATIONS SEE PROFILE		Somnath Bhowmick The Cyprus Institute 17 PUBLICATIONS 34 CITATIONS SEE PROFILE

Some of the authors of this publication are also working on these related projects:

Investigation of Low energy electron induced single strand break in DNA fragments View project

Advanced computing in complex systems physics, theoretical chemistry and stochastic models (IB16013) View project

Contents lists available at ScienceDirect



Computational and Theoretical Chemistry

journal homepage: www.elsevier.com/locate/comptc



Theoretical investigation of $X_{12}O_{12}$ (X = Be, Mg, and Ca) in sensing CH₂N₂: A DFT study



Mohsen Doust Mohammadi^a, Hewa Y. Abdullah^{b,*}, Somnath Bhowmick^c, George Biskos^{c,d}

^a School of Chemistry, College of Science, University of Tehran, Tehran 14176, Iran

^b Physics Education Department, Faculty of Education, Tishk International University, 44001 Erbil, Iraq

^c Climate & Atmosphere Research Centre, The Cyprus Institute, 20 K. Kavafi Street, Nicosia 2121, Cyprus

^d Faculty of Civil Engineering and Geosciences, Delft University of Technology, Delft 2628 CN, the Netherlands

ARTICLE INFO

Keywords: DFT Diazomethane Nanocage NBO QTAIM

ABSTRACT

The feasibility of detecting diazomethane (CH_2N_2) in the gas phase by adsorption onto the exterior surface of inorganic-based $X_{12}O_{12}$ (where X can be Be, Mg, or Ca) nanocages is investigated here using DFT. All the structures, including those of the pristine CH_2N_2 and of the nanocages, as well as of the CH_2N_2 /nanocage systems, have been optimized using the B3LYP-D3, M06-2X, ω B97XD, and CAM-B3LYP functionals, in conjunction with 6-311G(d) basis set. NBO, NCI, and QTAIM analyses results are in good agreement with each other. Furthermore, the Density Of States (DOSs), the natural charges, the Wiberg Bond Indices (WBI), and natural electron configurations were considered to investigate the nature of intermolecular interactions. The energy calculations indicate a strong size-dependent adsorption, with the nanocages comprised of large atoms being able to attract CH_2N_2 more strongly, and hence bind with it more effectively. The adsorption incurs also significant changes to HOMO and LUMO energies.

1. Introduction

Diazomethane (CH_2N_2) is a commonly used methylating agent that is a strong respiratory irritant, posing significant health concerns. At room temperature it is an extremely explosive gas, and thus comprises an important safety hazard [1]. As a result, there is an increasing demand in designing and developing sensors that can measure the concentration of diazomethane in the air for safety reasons.

Among the various types of sensors, those relying on the adsorption of the target gas (i.e., diazomethane in our case) on the sensing material, which consequently alter one measurable property (e.g., their resistance [2] are very promising for a number of reasons including their simplicity and low fabrication cost. Designing such sensing materials, however, requires a good understanding of the interaction (i.e., adsorption) between the target gas and the sensing element, which in turns warrants for systematic theoretical calculations.

Density functional theory (DFT), due to its high computational effectiveness for large molecules and its considerable accuracy, provides a suitable framework for theoretical studies of various chemical systems [3]. As a result, it is widely employed to understand the properties of chemical structures and material systems. At the same time, theoretical

studies on nanoscaled structures have become well-established in computational chemistry, providing numerous computational tools to study intermolecular interactions and to design materials that can exhibit enhanced performance for specific applications [4–11]. For instance, a wide range of nanomaterials have been developed and tested for gas sensors that can have low limits of detection, high measurement accuracy, low response time, resistant to variable environmental conditions, as well as low production cost [12–22].

Since the discovery of the C_{60} fullerene (which can be considered as the first synthesized nanocage) by Kroto et al. [23], numerous studies have investigated its unique properties, which distinguish it as a new carbon allotrope, in view of specific applications. Several experimental and theoretical studies carried out over the past decade have extended this activity by developing and studying structures similar to those of fullerenes, using elements from the third through the fifth groups of the periodic table [24–26]. Following this development, a significant number of studies have investigated the properties of these nanostructures for potential applications in medicine [27], optics [28], quantum computing [29], and electronics [30].

 X_nY_n (where X can be B, Al, or Ga, and Y can be N, P, or As) nanocages have a number of unique properties that can be useful for

* Corresponding author. *E-mail address:* hewayaseen@gmail.com (H.Y. Abdullah).

https://doi.org/10.1016/j.comptc.2021.113168

Received 5 November 2020; Received in revised form 15 January 2021; Accepted 18 January 2021 Available online 1 February 2021 2210-271X/© 2021 Elsevier B.V. All rights reserved. specific gas sensors. For instance, boron nitride (B_nN_n) nanocages are isoelectronic, which is in contrast to carbon fullerene-like nanocages, and exhibit unique optical, magnetic, and electronic properties. In this regard, the Coulomb blockade, photoluminescence and super magnetism of B_nN_n nanocages are significantly disparate compared to those of fullerenes [31]. Earlier theoretical investigations using DFT show the feasibility of adsorption of specific gaseous molecules, including formaldehyde [26], ammonia [32], CO [33], CO₂ [34], NO, N₂O [35], H₂, N₂ and CH₄ [36], *etc.*, on $B_{12}N_{12}$ nanocages. Similar theoretical studies, but using other nanocages, including Ga₁₂N₁₂ [37], $B_{12}P_{12}$ [25], Ga₁₂P₁₂ [38], $B_{12}As_{12}$ [39], Al₁₂As₁₂ [40], Ga₁₂As₁₂ [41], have also been reported.

Earlier studies have shown that $X_{12}O_{12}$ (X = Be, Mg, and Ca) nanocages can act as very effective adsorbents due to the highly ionic metal-oxide bond [42]. For example, Farrokhpour et al. have demonstrated that sulphur mustard can effectively adsorb on Mg12O12 nanocages [43], whereas Solimannejad et al. have theoretically shown that diazomethane can be adsorbed on silicon carbide nanotubes [44]. Here we report a systematic study on the adsorption of diazomethane molecule onto all six available adsorption sites of $X_{12}O_{12}$ (X = Be, Mg, and Ca) fullerene-like nanocages. The electronic properties, energetic parameters, and geometry optimization of all structures were investigated using Natural Bond Orbital (NBO), Non-Covalent Interaction (NCI), and Quantum Theory of Atoms In Molecules (QTAIM) analysis. All the geometry optimization calculations have been carried out by DFT using four different functionals (i.e., B3LYP-D3, M06-2X, ωB97XD, and CAM-B3LYP), together with a split-valence triple-zeta (6-311G(d)) basis set. The nature of the intermolecular interactions between the gas molecule and the above-mentioned nanocages were studied.

2. Computational details

All the structures investigated in this work have been optimized (assuming vacuum conditions) using the Kohn-Sham DFT [45,46]. Various functionals including CAM-B3LYP [47], M06-2X [48], ω B97XD [49], and the B3LYP-D3 scheme of Grimme [50], together with splitvalence triple-zeta basis sets with d-type Cartesian-Gaussian polarization functions (6-311G(d)) were employed. It should be noted that the 6-311G(d) basis set is suitable for the systems considered in this work, as reported in a benchmark study by Grimme et al. [51]. No symmetry constraints were imposed. All the molecular geometries were built using the GaussView 6.0.16 [52] software package, which is fully optimized according to the Berny [53] method in Gaussian 16 Rev. C.01 [54] Linux based version of the software. The optimization processes were performed using the default Gaussian convergence criteria. A frequency calculation has been performed to ascertain the stability of each structure and to determine the nature of each stationary point. In addition, wave function stability calculations have also been used to evaluate the stability of the electrons. NBO calculations were carried out using the NBO v 3.1 software (which is integrated within the Gaussian package) to determine the charge-transfer interactions between occupied and unoccupied orbitals. To extract the result data of NBO, NCI, and QTAIM analysis, as well as to depict the DOS diagrams, we employed respectively the Multiwfn [55] and the GaussSum [56] software.

The energy of adsorption (E_{ads}) between two fragments (i.e., the nanocage and the CH₂N₂ gas molecule) can be considered as follows:

$$E_{ads} = E_{cage/gas} - E_{cage} - E_{gas} + \Delta E_{(BSSE)}$$
(1)

where $E_{cage/gas}$ is the total energy of the gas/nanocage cluster, E_{cage} and E_{gas} are the energies of the isolated nanocage and isolated gas molecule, respectively, and $\Delta E_{(BSSE)}$ is the Basis Set Superposition Error (BSSE) obtained from the Boys and Bernardi's counterpoise procedure [57] as follows:

$$\Delta E_{(BSSE)} = \Delta E_{cluster} - \Delta E_{cage}^{cluster} - \Delta E_{gas}^{cluster}$$
⁽²⁾



Fig. 1. The structure of the $X_{12}Y_{12}$ (where X is Be, Mg, or Ca and Y is O) nanocages, displaying all adsorption sites, including the X-site (on top of the X atom), Y-site (on top of the Y atom), H-6 site (on top of the hexagonal ring), H-4 site (on top of the tetragonal ring), XY66 site (on top of the atomic bond between two hexagonal rings) and XY64 site (on top of the atomic bond between tetragonal and hexagonal rings).

According to Equation (1), negative values of E_{ads} (i.e., corresponding to exothermic adsorptions) indicate that the formed complex is stable; otherwise, positive values of E_{ads} belong to a local minimum in which the adsorption of CH₂N₂ on the surface of the nanocage is deterred by an energy barrier.

3. Results and discussion

3.1. Structural analysis

The structures of pristine $X_{12}O_{12}$ nanocages were optimized using DFT at the CAM-B3LYP/6-311G(d) level. Although CAM-B3LYP describes the long-range interactions, it does not consider dispersion correction and the charge transfer excitations; therefore, all of the structures were re-optimized using robust methods (i.e., M06-2X, ω B97XD, and B3LYP-D3). Head-Gordon et al. [49] developed the ω B97XD functional to account to capture the effect of long-range interactions and to make dispersion-related corrections. Among the functionals of the Minnesota 06 suit, developed by Truhlar et al. [48], the global hybrid functional with double non-local exchange (2X) amounts (i.e., M06-2X) is a high-performance method to study the non-covalent interactions. To take into account the effect of dispersion, the latest version of B3LYP-D3, known as D3 (BD) (GD3BJ) developed by Grimme et al. [50], would also be a good option and has been used in this present study.

The geometrical structure of the $X_{12}Y_{12}$ nanocages containing the X-Y bonds is shown in Fig. 1. These nanocages consist of eight (8) symmetric hexagons and six (6) tetragons having an equal number of adsorption positions for the adsorption of any molecule onto their outer surface shown further below. The adsorption sites can be categorized as follows: on top of the X atom (T_1), on top of the Y atom (T_2), on top of the XY66 bond (T_3), on top of the H-6 position (T_4), on top of the XY64 bond (T_5), and on top of H-4 position (T_6). After the optimization of the pristine nanocage structures, we placed the CH₂N₂ at a vertical distance of approximately 2.5 Å onto the outer surface of the nanocages in any of the six possible adsorption sites.

The optimized geometrical parameters of the CH_2N_2 molecule obtained by all four functionals are in good agreement with experimental results (see Fig. S1 in the supplemental material). Evidently, the B3LYP-D3 method gives the best results among all four functionals compared to the observed fundamental vibrational frequencies (see Table S1 in the supplemental material). For example, the B3LYP-D3 C—H stretching





Fig. 2. Optimized structure of CH_2N_2 gas molecule obtained from B3LYP-D3/6-311G(d) level of theory. The unit of the bond lengths are given in Å and the angles in degrees.

frequency was calculated to be 3179 cm^{-1} , which is in good agreement with the experimental value of 3073 cm^{-1} . The optimized structural parameters of the isolated CH₂N₂ gas molecule obtained by the B3LYP-D3 method is depicted in Fig. 2.

The optimized structures of the bare $X_{12}O_{12}$ nanocages calculated by the functionals considered in this study are provided in the supplementary material. All four methods have good correspondence with the previous B3LYP/6-31+G(d) calculations by Omidi et al. [42]. For example, the X-O bond lengths calculated in this study lies between 1.58 and 1.61 Å when X is Be, 1.93–1.95 Å when X is Mg and 2.18–2.19 Å when X is Ca, having good agreement with the values obtained by obtained by Omidi et al.: i.e., 1.58 Å when X is Be, 1.95 Å when X is Mg and 2.28 Å when X is Ca. The final B3LYP-D3 optimized structures of the pristine nanocages together with the gas/cage complexes and the corresponding bond lengths are illustrated in Figs. 3 and 4 and the corresponding atomic co-ordinates are provided in the supplementary material.

It should be stressed that to find the optimum distance between the nanocage and the adsorbed gas molecule, we used rigid scans in the





Fig. 3. The bond length (Å) obtained at B3LYP-D3/6-311G(d) level for (a) $Be_{12}O_{12}$, (b) $Mg_{12}O_{12}$, and (c) $Ca_{12}O_{12}$.



Fig. 4. The optimized structures of (a) CH₂N₂/Be₁₂O₁₂, (b) CH₂N₂/Mg₁₂O₁₂, and (c) CH₂N₂/Ca₁₂O₁₂ obtained from B3LYP-D3/6-311G(d) level of theory. The units of the bond lengths are given in Å.

Table 1

Adsorption energies (E_{ads}) for CH₂N₂/X₁₂O₁₂ (where X is Be, Mg, or Ca) complexes calculated at X, Y, H-4, H-6, XY64 and XY66 adsorption sites. All values are expressed in eV and are obtained from geometry optimization calculations using the B3LYP-D3 functional in combination with 6-311G(d) basis set.

System	Х	Y	H-4	H-6	XY64	XY66
CH ₂ N ₂ /Be ₁₂ O ₁₂	0.012	0.014	0.002	0.030	0.0	0.014
CH ₂ N ₂ /Mg ₁₂ O ₁₂	0.009	0.001	0.002	0.004	0.0	0.012
CH ₂ N ₂ /Ca ₁₂ O ₁₂	0.005	0.015	0.003	0.001	0.0	0.008

PM7/6-311G(d) level of theory. The logical approach is to place the gas molecule in each of these positions and calculate the amount of adsorption energy (E_{ads}). Our experience shows that negligible differences exist in the amounts of adsorption energies when we place the gas in any of the six possible adsorption sites. As mentioned by Foresman et al. [58], when the differences in the adsorption energies are "below the range of chemical interest", placing the gas molecule in different positions on the nanocages provides identical results. Next, regarding the amount of E_{ads} values, some structures were chosen for DFT calculations at CAM-B3LYP/6-311G(d) level of theory. Finally, structures with the most negative energy were selected for the re-optimization process using the M06-2X, ωB97XD, and B3LYP-D3 functional and the 6-311G(d) basis set. The B3LYP-D3 functional is the basis for decision making to study interactions that consider dispersion corrections and

Table 2

Adsorption energies (E_{ads}) for CH₂N₂/X₁₂O₁₂ (where X is Be, Mg, or Ca) complexes at the XY64 adsorption site. All values are expressed in eV and are obtained from geometry optimization calculations using the CAM-B3LYP, M06-2X, ω B97XD, and B3LYP-D3 functionals in combination with 6-311G(d) basis set.

Systems	B3LYP-D3	CAM-B3LYP	M06-2X	ωB97XD
- ,				
CH2N2/Be12O12	-0.712	-0.605	-0.804	-0.676
CH ₂ N ₂ /Mg ₁₂ O ₁₂	-0.743	-0.546	-0.748	-0.633
$CH_2N_2/Ca_{12}O_{12}$	-0.699	-0.446	-0.610	-0.521

are in good agreement with the experimental observations, whereas the rest of the functionals are given here for comparison only. The E_{ads} values obtained from optimization calculations are provides in Tables 1 and 2.

The adsorption energies (E_{ads}) calculated by the B3LYP-D3 method at all six adsorption sites are reported in Table 1. As discussed previously, the choice of the adsorption site has a negligible effect on the corresponding adsorption energies, although the XY64 site releases the largest energy compared to others. The maximum difference in E_{ads} among all adsorption sites is found to be around 0.03 eV. Table 2 provides the calculated E_{ads} on the $X_{12}O_{12}$ nanocages at the XY64 site obtained using different functionals. From these values, it is clear that as the size of the atoms in the nanocages decreases, the intensity of adsorption between the gas molecule and the nanocages increases. For

M.D. Mohammadi et al.

Table 3

HOMO energy ($\epsilon_{\rm H}$), LUMO energy ($\epsilon_{\rm L}$), Fermi energy ($E_{\rm F}$), HOMO–LUMO energy gap (HLG), chemical potential (μ), chemical hardness (η), and electrophilicity (ω) of the pristine X₁₂O₁₂ nanocages and the CH₂N₂/X₁₂O₁₂ complexes (where X is Be, Mg, or Ca). All values are in eV and were obtained using the B3LYP-D3/6-311G(d) level of theory.

Systems	$\varepsilon_{ m H}$	$\varepsilon_{\rm L}$	E_F	HLG	μ	η	ω
Be ₁₂ O ₁₂	-8.672	-0.237	-4.455	8.435	-4.455	4.218	41.848
Mg ₁₂ O ₁₂	-6.543	-1.680	-4.111	4.864	-4.111	2.432	20.553
Ca ₁₂ O ₁₂	-4.649	-1.089	-2.869	3.560	-2.869	1.780	7.324
CH2N2/Be12O12	-7.723	-2.393	-5.058	5.330	-5.058	2.665	34.090
CH ₂ N ₂ /Mg ₁₂ O ₁₂	-6.287	-2.091	-4.189	4.196	-4.189	2.098	18.406
CH2N2/Ca12O12	-4.527	-1.931	-3.229	2.596	-3.229	1.298	6.766



Fig. 5. Density of state maps for **(a)** $Be_{12}O_{12}$, **(b)** $CH_2N_2/Be_{12}O_{12}$, **(c)** $Mg_{12}O_{12}$, **(d)** $CH_2N_2/Mg_{12}O_{12}$, **(e)** $Ca_{12}O_{12}$, and **(f)** $CH_2N_2/Ca_{12}O_{12}$. Data were obtained from the B3LYP-D3/6-311G(d) level of theory.

Table 4

Values of Mulliken, Mayer, and Wiberg, Fuzzy, and Laplacian bond orders obtained for the interactions of the CH_2N_2 molecule and $X_{12}O_{12}$ (where X is Be, Mg, or Ca). All calculations were performed using the B3LYP-D3/6-311G(d) level of theory.

System	$X^a {\cdots} Y^b$	Mulliken	Mayer	FBO	WBI	Laplacian
CH ₂ N ₂ /Be ₁₂ O ₁₂	C…Be	0.204	0.259	0.449	0.528	0.112
CH ₂ N ₂ /Mg ₁₂ O ₁₂	C…Mg	0.203	0.214	0.442	0.391	0.084
CH ₂ N ₂ /Ca ₁₂ O ₁₂	C…Ca	0.099	0.128	0.252	0.180	0.069

^a X atoms are belong to CH₂N₂.

^b Y atoms are belong to Nanocages.

example, in the case of CAM-B3LYP/6-311G(d) optimization of $CH_2N_2/Be_{12}O_{12}$, $CH_2N_2/Mg_{12}O_{12}$, and $CH_2N_2/Ca_{12}O_{12}$, the adsorption energies are -0.605, -0.546, and -0.446 eV, respectively. It should be noted here that the B3LYP-D3 adsorption energy for diazomethane obtained in this work (-0.71 eV to -0.79 eV) is comparable to the adsorption of sulfur mustard on $Be_{12}O_{12}$ and $Mg_{12}O_{12}$ nanocages (-0.47 eV to -0.99 eV) [43]. This is despite the fact that here we used a triple zeta basis set, in contrast to a double zeta basis that has been employed for the sulfur mustard adsorption study.

3.2. Electronic structure

The "Conceptual DFT" has been developed to consider the reactivity concept. Various properties can be obtained from the HOMO-LUMO energy gap (HLG) [59]. For example, when an external potential applied to a system, within the Hohenberg-Kohn context [41,42], the HOMO and LUMO energies are related to the ionization potential (IP) and electron affinity (EA). One can also easily show that the electronegativity (χ) is equal to the negative chemical potential (μ). Furthermore, the electrophilicity index (ω) is inversely related to the chemical hardness (η), which in turn is related to the difference between IP and EA. The calculated values of these properties are listed in Table 3.

The HOMO-LUMO energy gap (ΔE_g) of Be₁₂O₁₂, Mg₁₂O₁₂, and Ca₁₂O₁₂ have been calculated to be 8.43, 4.86, and 3.56 eV, respectively, at the B3LYP-D3/6-311G(d) theory level. The adsorption of CH₂N₂ on its surface will reduce the energy gap to 5.33, 4.20, and 2.60 eV, respectively. These results show the sensitivity of the HOMO-LUMO energies with respect to adsorption. From the results provided in Table 3, it is also evident that the LUMO values were largely stabilized compared to the destabilization of the HOMO orbitals.

The sensor response (*S*) is defined as $S = exp\left(\frac{|\Delta E_g|}{kT}\right) - 1$, where *k* is

the Boltzmann constant and *T* the temperature expressed in Kelvin. It is clear that the resistivity diminishes when the energy gap ΔE_g is reduced since the resistivity is proportional to the reciprocal of the conducting electron population. Hence, the resistivity of the tube upon adsorption of gas molecules is low, and the electric current generated in the circuit will exhibit the lowest resistance. The density of state (DOS) map is useful for intuitively revealing the density of distribution of the molecular orbitals in different energy regions, and the gap is directly visible from this map (see Fig. 5).

3.3. Natural Bond Orbital (NBO) analysis

The NBO method, developed by Weinhold et al. [60], is one of the most robust population analysis methods that can determine the electron density distribution within bonds between the atoms. The term NBO refers to a bonding orbital with the maximum electron density. A density matrix, calculated from the DFT method, and the atomic charge, is used to define the natural bonding orbitals. In order to complete the span of the valance space, in addition to the bonding NBOs (σ), determination of antibonding NBOs (σ^*) is also required. In the present study, the NBO calculations have been performed to evaluate various types of

Table 5

Natural charge (in *a.u.*) and natural electron configurations for the CH_2N_2 and $CH_2N_2/X_{12}O_{12}$ structures (where X is Be, Mg, or Ca). All values have been obtained from NBO analysis using the B3LYP-D3/6-311G(d) level of theory.

Systems	atom	Natural Charge	Natural Electron Configuration
CH ₂ N ₂	C H1 H2 N1 N2	-0.480 0.243 0.243 0.021 -0.026	[core] 2 s(1.03) 2p(3.43) 3p(0.01) 1s(0.76) 1s(0.76) [core] 2s(1.14) 2p(3.81) 3s(0.01) 3p (0.02) 3d(0.01) [core] 2s(1.61) 2p(3.39) 3s(0.01) 3p
CH ₂ N ₂ / Be ₁₂ O ₁₂	С	-0.728	(0.01) 3 <i>d</i> (0.01) [core] 2 <i>s</i> (1.17) 2 <i>p</i> (3.53) 3 <i>s</i> (0.01) 3 <i>p</i> (0.02)
	H1	0.279	1s(0.72)
	H2	0.279	1s(0.72)
	N1	0.107	[core] 2s(1.16) 2p(3.71) 3s(0.01) 3p (0.02)
	N2	0.117	[core] 2s(1.60) 2p(3.25) 3s(0.01) 3p (0.01) 3d(0.02)
CH ₂ N ₂ / Mg ₁₂ O ₁₂	С	-0.661	[core] 2s(1.12) 2p(3.50) 3s(0.01) 3p (0.02)
012 12	H1	0.271	1s(0.73)
	H2	0.271	1s(0.73)
	N1	0.089	[core] 2s(1.15) 2p(3.73) 3s(0.01) 3p (0.02) 3d(0.01)
	N2	0.064	[core] 2s(1.60) 2p(3.31) 3s(0.01) 3p (0.01) 3d(0.01)
$CH_2N_2/Ca_{12}O_{12}$	С	-0.557	[core] 2s(1.07) 2p(3.46) 3p(0.02)
12 - 12	H1	0.261	1s(0.74)
	H2	0.261	1s(0.74)
	N1	0.077	[core] $2s(1.14) 2p(3.75) 3s(0.01) 3p$ (0.02) $3d(0.01)$
	N2	-0.038	[core] $2s(1.60) 2p(3.41) 3s(0.01) 3p$ (0.01) $3d(0.01)$

bond order, including Mulliken [61] and Mayer [62] bond order as well as Wiberg bond index (WBI) in Löwdin orthogonalized basis [63]. Mulliken and Mayer's bond orders are sensitive to the chosen basis set, especially for the basis set that includes diffuse functions. On the other hand, the Wiberg bond order is less dependent on the basis set in comparison to the other two. Additionally, the Fuzzy Bond Order (FBO) was proposed by Mayer [64], and the Laplacian bond order proposed by Lu [65] has also been calculated from the corresponding overlap and density matrix. In Table 4, we report the calculated C-X bond orders based on these different methods. Evidently, there is a correlation between the size of the cation in the nanocages and the bond order: *i.e.*, the larger the cation, the lower the bond order. The bond order values also signify that the interaction of the gas with the nanocages can be classified as strong interactions.

We have also calculated the natural charges and natural electron configuration data through natural population analysis combined with NBO (cf. Table 5). Using the NBO analysis, the nature of the intermolecular interaction between the gas molecule and nanocages becomes more intelligible. As reflected by the calculates reported in Table 5, the existence of charge transfer within these complexes is very obvious. The valance shell of each atom in CH₂N₂ is changed due to the interaction with the nanocages. For example, the valence shell of C atom in molecules of the gas phase is 2s(1.03) 2p(3.43) 3p(0.01), which is changed to 2s(1.17) 2p(3.53) 3s(0.01) 3p(0.02) for the Be₁₂O₁₂/gas complex during the adsorption process. This result suggests that a strong interaction has occurred.

3.4. Quantum Theory of Atoms In Molecules (QTAIM) analysis

QTAIM analysis has been used to study the bond types and intermolecular interactions. A critical point of the electron density, including

Table 6

QTAIM topological parameters for electron density $\rho(\mathbf{r})$, Laplacian of electron density $\nabla^2 \rho(\mathbf{r})$, kinetic electron density $G(\mathbf{r})$, potential electron density $V(\mathbf{r})$, eigenvalues of Hessian matrix (λ), and bond ellipticity index (ε) at the BCPs of the CH₂N₂ clusters with X₁₂O₁₂ (where X can be Be, Mg, or Ca). All values were calculated using the B3LYP-D3/6-311G(d) level of theory and NBO analysis.

Systems	Bond	ρ	$\nabla^2 r$	<i>G</i> (r)	V(r)	$G(\mathbf{r})/ V (\mathbf{r})$	λ_1	λ_2	λ_3	ε
CH2N2/Be12O12	С…Ве	0.0462	0.2009	0.0561	-0.0621	0.9045	-0.0640	-0.0682	0.3331	0.0671
CH ₂ N ₂ /Mg ₁₂ O ₁₂	C…Mg	0.0218	0.1029	0.0228	-0.0200	1.1443	-0.0231	-0.0232	0.1492	0.0013
CH2N2/Ca12O12	C…Ca	0.0130	0.0468	0.0102	-0.0086	1.1791	-0.0101	-0.0109	0.0679	0.0823
	N…Ca	0.0119	0.0484	0.0104	-0.0087	1.1963	-0.0101	-0.0103	0.0688	0.0133



Fig. 6. The bond critical points graph for the (a) CH₂N₂/Be₁₂O₁₂, (b) CH₂N₂/Mg₁₂O₁₂, and (c) CH₂N₂/Ca₁₂O₁₂systems. The orange dots represent the BCPs.

minimum, maximum, or saddle point, can belong to: (1) *atomic critical point* (ACP); (2) *bond critical point* (BCP); (3) *ring critical point* (RCP); and (4) *cage critical point* (CCP). The electron density $\rho(\mathbf{r})$ and the Laplacian electron density $\nabla^2 \rho(\mathbf{r})$ play an essential role in the QTAIM analysis since they determine the segmentation and identification of different types of chemical interactions. A bond critical point with negative values of the Laplacian electron density and large values of the electron density $(\rho(\mathbf{r}) > 10^{-1} a.u.)$ is defined as a covalent bond. On the other hand, the positive value of $\nabla^2 \rho(\mathbf{r})$ designates that the interactions can be classified as of the non-substrate close-shell type (which include ionic and van der Waals interactions) [66].

The values of Lagrangian kinetic energy $G(\mathbf{r})$ and potential energy density $V(\mathbf{r})$ divulges the nature of the intermolecular interaction; therefore, the ratio of $G(\mathbf{r})/|V|(\mathbf{r})$ can be hired as an appropriate index in link classification. When $G(\mathbf{r})/|V|(\mathbf{r}) < 0.5$, the nature of the interaction is covalent, and if $G(\mathbf{r})/|V|(\mathbf{r}) > 1.0$, the interaction is non-covalent. On the other hand, a large value of the elliptical bond (ε) represents an unstable structure and is related to the ratio between the eigenvalues (λ_1 , λ_2 ,... etc.) of the Hessian matrix [67].

Table 6 provides the calculated values of these parameters obtained from the QTAIM analysis. Since all values for $\nabla^2 \rho(\mathbf{r})$ are positive and the $\rho(\mathbf{r})$ values are always < 0.1, we can safely assume that the stabilities of the adsorbed complexes originate from the non-covalent or van der Waals interactions. Moreover, the $G(\mathbf{r})/|V|(\mathbf{r})$ ratios are higher than unity, except for the C…Be interaction in the CH₂N₂/Be₁₂O₁₂ cluster. These results suggest that intermolecular interactions can be classified as non-covalent. The calculated values of the elliptical bonds are also very small, indicating that the structures are highly stable. Fig. 6 provides a schematic illustration of the bond critical points between gas and nanocages.

The results of QTAIM analysis discussed in the previous section showed that the interactions between CH_2N_2 and nanotubes are essentially non-covalent; hence, it is useful to re-evaluate the interaction with a non-covalent analysis. The reduced density gradient (RDG) and

sign $\lambda_2(\mathbf{r})\rho(\mathbf{r})$ are a pair of functions used in non-covalent interaction (NCI) [68] analysis, which can be implemented to visualize the region and the type of weak interactions. Plots of these two functions, RDG and sign $\lambda_2(\mathbf{r})\rho(\mathbf{r})$, can be used to identify specific areas; in particular, the non-covalent interaction area. The points that indicate strong interactions are located in the sign $\lambda_2(\mathbf{r})\rho(\mathbf{r}) < 0$ region. Relatively weak van der Waals interactions are found in the sign $\lambda_2(\mathbf{r})\rho(\mathbf{r}) \approx 0$ region. Furthermore, if points are in sign $\lambda_2(\mathbf{r})\rho(\mathbf{r}) > 0$ region, suggests that the interactions are repulsive [68,69]. As it turns out, bond strength is closely related to the density matrix $\rho(\mathbf{r})$ as well as to sign λ_2 . Low RDG and low electron density regions should be further consulted to determine whether non-covalent interactions occur between the two components involved in the adsorption process.

From Fig. 7, it is evident that non-covalent interaction analysis can adequately describe the interactions between gas and nanocages. In the sign $\lambda_2(\mathbf{r})\rho(\mathbf{r}) < 0$ region, many points are observed, which indicates that there are very strong non-covalent interactions between the two fragments. It is also observed that in the sign $\lambda_2(\mathbf{r})\rho(\mathbf{r}) \approx 0$ region, there is no high concentration of points. Therefore, it can be concluded that the intermolecular interactions during the adsorption process are not van der Waals.

4. Conclusion

The intermolecular interactions between the CH_2N_2 gas molecule and $X_{12}O_{12}$ (where X can be Be, Mg, or Ca) nanocages were studied by DFT. All molecular structures have been optimized using the CAM-B3LYP, M06-2X, ω B97XD, and B3LYP-D3 functionals together with a 6-311G(d) basis set. The optimized structures obtained from B3LYP-D3/ 6-311G(d) were chosen for population analysis calculations because of their good agreement with earlier experimental and theoretical results. The calculated values of the adsorption energy indicate that on top of the X-O bond connecting the tetragonal and hexagonal rings is the most favourable adsorption site. In general, adsorption energy increases with



Fig. 7. The sign $(\lambda_2)\rho(\mathbf{r})$ (X-axis) vs. RDG (Y-axis) diagrams for (a) Be₁₂O₁₂, (b) CH₂N₂/Be₁₂O₁₂, (c) Mg₁₂O₁₂, (d) CH₂N₂/Mg₁₂O₁₂, (e) Ca₁₂O₁₂, and (f) CH₂N₂/Ca₁₂O₁₂. The diagrams were obtained from B3LYP-D3/6-311G(d) level of theory.

the size of the central metal atom of the $X_{12}O_{12}$ nanocages. Several bond order analysis data corroborate the above conclusion. To understand the nature of the intermolecular interactions, we further performed NBO, QTAIM, and NCI analyses, with the results appearing to correspond well with each other. The NBO calculations show a significant charge transfer, whereas the NCI and QTAIM results indicate strong intermolecular interactions. To sum up, we conclude that the studied $X_{12}O_{12}$ nanocages have an active surface in favor of the CH₂N₂ gas adsorption. Our findings indicate that nanomaterials containing $X_{12}O_{12}$ nanocages investigated here would be suitable for designing sensitive CH₂N₂ sensors.

CRediT authorship contribution statement

Mohsen Doust Mohammadi: Writing - original draft, formal analysis and review & editing. Hewa Y. Abdullah: Supervision, Investigation and project administration. Somnath Bhowmick: Conceptualization, and validation. George Biskos: Resources, and visualization.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

HYA thanks the Solid-State Theory Group, Physics Department, Università degli Studi di Milano, Milan, Italy for providing computational facilities. SB and GB acknowledge the support of the European Regional Development Fund and the Republic of Cyprus through the Research Innovation Foundation (Cy-Tera project under the grant NEA YPODOMH/STPATH/0308/31, and NANO²LAB project under the grant INFRASTRUCTURES/1216/0070).

Appendix A. Supplementary material

Supplementary data to this article can be found online at https://doi.org/10.1016/j.comptc.2021.113168.

References

- D. Dallinger, C.O. Kappe, Lab-scale production of anhydrous diazomethane using membrane separation technology, Nat. Protoc. 12 (2017) 2138–2147.
- [2] N.A. Isaac, M. Valenti, A. Schmidt-Ott, G. Biskos, Characterization of Tungsten Oxide Thin Films Produced by Spark Ablation for NO₂ Gas Sensing, ACS Appl. Mater. Interfaces 8 (2016) 3933–3939.
- [3] D. Sholl, J.A. Steckel, Density Functional Theory: A Practical Introduction, John Wiley & Sons, Inc., Hoboken, New Jersey, 2009.
- [4] A.S. Novikov, M.L. Kuznetsov, Theoretical study of Re(IV) and Ru(II) bisisocyanide complexes and their reactivity in cycloaddition reactions with nitrones, Inorg. Chim. Acta 380 (2012) 78–89.
- [5] A.S. Novikov, M.L. Kuznetsov, A.J.L. Pombeiro, Theory of the formation and decomposition of n-heterocyclic aminooxycarbenes through metal-assisted [2+3]dipolar cycloaddition/retro-cycloaddition, Chem. A Eur. J. 19 (2013) 2874–2888.
- [6] A.S. Novikov, M.L. Kuznetsov, A.J.L. Pombeiro, N.A. Bokach, G.B. Shul'pin, Generation of HO• radical from hydrogen peroxide catalyzed by aqua complexes of the Group III metals [M(H2O)n]3+ (M = Ga, In, Sc, Y, or La): A Theoretical Study, ACS Catal. 33 (2013) 1195–1208.
- [7] A.A. Melekhova, A.S. Novikov, K.V. Luzyanin, N.A. Bokach, G.L. Starova, V. V. Gurzhiy, V.Y. Kukushkin, Tris-isocyanide copper(I) complexes: Synthetic, structural, and theoretical study, Inorg. Chim. Acta 434 (2015) 31–36.
- [8] A.S. Novikov, 1,3-Dipolar cycloaddition of nitrones to transition metal-bound isocyanides: DFT and HSAB principle theoretical model together with analysis of vibrational spectra, J. Organomet. Chem. 797 (2015) 8–12.
- [9] K.I. Kulish, A.S. Novikov, P.M. Tolstoy, D.S. Bolotin, N.A. Bokach, A.A. Zolotarev, V.Y. Kukushkin, Solid state and dynamic solution structures of O-carbamidine amidoximes gives further insight into the mechanism of zinc(II)-mediated generation of 1,2,4-oxadiazoles, J. Mol. Struct. 1111 (2016) 142–150.
- [10] A.S. Novikov, M.L. Kuznetsov, B.G.M. Rocha, A.J.L. Pombeiro, G.B. Shul'pin, Oxidation of olefins with H₂O₂ catalysed by salts of group III metals (Ga, In, Sc, Y and La): epoxidation versus hydroperoxidation, Catal. Sci. Technol. 6 (2016) 1343–1356.
- [11] F. Li, H. Asadi, DFT study of the effect of platinum on the H₂ gas sensing performance of ZnO nanotube: explaining the experimental observations, J. Mol. Liq. (2020) 113139.
- [12] M.D. Mohammadi, H.Y. Abdullah, The adsorption of chlorofluoromethane on pristine, and Al-and Ga-doped boron nitride nanosheets: a DFT, NBO, and QTAIM study, J. Mol. Model. 26 (2020) 1–15.
- [13] M.D. Mohammadi, H.Y. Abdullah, Theoretical study of the adsorption of amantadine on pristine, Al-, Ga-, P-, and As-doped boron nitride nanosheets: a PBC-DFT, NBO, and QTAIM study, Theor. Chem. Acc. 139 (2020) 1–17.
- [14] M.D. Mohammadi, H.Y. Abdullah, The adsorption of bromochlorodifluoromethane on pristine and Ge-doped silicon carbide nanotube: a PBC-DFT, NBO, and QTAIM study, Struct. Chem. 32 (2021) 481–494.
- [15] M.D. Mohammadi, I.H. Salih, H.Y. Abdullah, An Ultimate Investigation on the adsorption of amantadine on pristine and decorated fullerenes C59X (X= Si, Ge, B, Al, Ga, N, P, and As): A DFT, NBO, and QTAIM Study, J. Theor. Comput. Chem. (2020) 1–17.
- [16] Mohsen Doust Mohammadi, H.Y. Abdullah, The adsorption of Bromochlorodifluoromethane on Pristine, Al, Ga, P, and As-doped Boron Nitride Nanotubes: A Study Involving PBC-DFT, NBO Analysis, and QTAIM, Comput. Theoret. Chem. 1193 (2021) 113047.
- [17] Mohsen Doust Mohammadi, Idris H. Salih and H. Y. Abdullah The adsorption of chlorofluoromethane on pristine and Ge-doped silicon carbide nanotube: a PBC-DFT, NBO, and QTAIM study molecular simulation, 46 (2020)1405–1416.
- [18] M.D. Mohammadi, H.Y. Abdullah, The Adsorption of Chlorofluoromethane on Pristine, Al, Ga, P, and As-doped Boron Nitride Nanotubes: A PBC-DFT, NBO, and QTAIM Study (2020) 12115–12124.
- [19] E. Nemati-Kande, M. Abbasi, M. Doust Mohammadi, DFT, QTAIM and NBO Investigation of the Interaction of Rare Gases with Pristine and Decorated Boron Nitride Nanotube, Chem. Select 3 (2018) 9833–9840.

- [20] M. Nemati-Kande, M.D. Abbasi, Mohammadi, DFT studies on the interactions of pristine, Al and Ga-doped boron nitride nanosheets with CH3X (X= F, Cl and Br), J. Mol. Struct. 1199 (2020), 126962.
- [21] E. Nemati-Kande, M. Abbasi, M.D. Mohammadi, Feasibility of Pristine and Decorated AlN and SiC Nanotubes in Sensing of Noble Gases: A DFT study, ChemistrySelect 4 (2019) 2453–2462.
- [22] M.D. Mohammadi, M. Hamzehloo, The adsorption of bromomethane onto the exterior surface of aluminum nitride, boron nitride, carbon, and silicon carbide nanotubes: A PBC-DFT, NBO, and QTAIM study, Comput. Theor. Chem. 1144 (2018) 26–37.
- [23] H.W. Kroto, J.R. Heath, S.C. O'Brien, R.F. Curl, R.E. Smalley, C60: Buckminsterfullerene, Nature 318 (1985) 162–163.
- [24] H.-S. Wu, F.-Q. Zhang, X.-H. Xu, C.-J. Zhang, H. Jiao, Geometric and energetic aspects of aluminum nitride cages, J. Phys. Chem. A 107 (2003) 204–209.
- [25] J. Beheshtian, Z. Bagheri, M. Kamfiroozi, A. Ahmadi, A comparative study on the B₁₂N₁₂, Al₁₂N₁₂, B₁₂P₁₂ and Al₁₂P₁₂ fullerene-like cages, J. Mol. Model. 18 (2012) 2653–2658.
- [26] T. Oku, A. Nishiwaki, I. Narita, Formation and atomic structure of B₁₂N₁₂ nanocage clusters studied by mass spectrometry and cluster calculation, Sci. Technol. Adv. Mater. 5 (2004) 635.
- [27] X. Michalet, F.F. Pinaud, L.A. Bentolila, J.M. Tsay, S. Doose, J.J. Li, G. Sundaresan, A. Wu, S. Gambhir, S. Weiss, Quantum dots for live cells, in vivo imaging, and diagnostics, Science 307 (2005) 538–544.
- [28] C. Yen, M. Mahmoud, M. El-Sayed, Photocatalysis in gold nanocage nanoreactors, J. Phys. Chem. A 113 (2009) 4340–4345.
- [29] K. Matsuura, N. Kimizuka, T. Yamashita, DNA nanocage by self-organization of DNA and process for producing the same, and DNA nanotube and molecule carrier using the same, Google Patents, 2008.
- [30] F. Meng, B. Sana, Y. Li, Y. Liu, S. Lim, X. Chen, Bioengineered tunable memristor based on protein nanocage, Small 10 (2014) 277–283.
- [31] H. Tavakol, D. Shahabi, DFT, QTAIM, and NBO study of adsorption of rare gases into and on the surface of sulfur-doped, single-wall carbon nanotubes, J. Phys. Chem. C 119 (2015) 6502–6510.
- [32] E. Vessally, E. Ahmadi, M.D. Esrafili, A. Hosseinian, Adsorption and decomposition of formaldehyde on the B₁₂N₁₂ nanostructure: a density functional theory study, Monatshefte für Chemie-Chemical Monthly 148 (2017) 1727–1731.
- [33] A.A. Peyghan, H. Soleymanabadi, Computational study on ammonia adsorption on the X₁₂Y₁₂ nano-clusters (X= B, Al and Y= N, P), Curr. Sci. (2015) 1910–1914.
- [34] J. Beheshtian, Z. Bagheri, M. Kamfiroozi, A. Ahmadi, Toxic CO detection by B₁₂N₁₂ nanocluster, Microelectron. J. 42 (2011) 1400–1403.
- [35] D. Quiñonero, A. Frontera, P.M. Deya, Feasibility of single-walled carbon nanotubes as materials for CO₂ adsorption: a DFT study, J. Phys. Chem. C 116 (2012) 21083–21092.
- [36] M.T. Baei, Si-Doped B₁₂N₁₂ Nanocage as an Adsorbent for Dissociation of N₂O to N₂ Molecule, Heteroat. Chem. 24 (2013) 476–481.
- [37] H. Cui, Y. Yong, H. Jiang, L. Yang, S. Wang, G. Zhang, M. Guo, X. Li, The M₁₂N₁₂ (M= Al, Ga) clusters as potential sensors for NO, NO₂ and HCN detection, Mater. Res. Express 4 (2017), 015009.
- [38] V. Tozzini, F. Buda, A. Fasolino, Spontaneous formation and stability of small GaP fullerenes, Phys. Rev. Lett. 85 (2000) 4554.
- [39] A. Ektarawong, S.I. Simak, B. Alling, First-principles prediction of stabilities and instabilities of compounds and alloys in the ternary B-As-P system, Phys. Rev. B 96 (2017), 024202.
- [40] L. Guo, Evolution of the electronic structure and properties of neutral and charged aluminum arsenide clusters: A comprehensive analysis, Comput. Mater. Sci. 42 (2008) 489–496.
- [41] J. Wang, L. Ma, J. Zhao, G. Wang, X. Chen, R. Bruce King, Electronic and magnetic properties of manganese and iron-doped Ga n As n nanocages (n=7–12), J. Chem. Phys. 129 (2008), 044908.
- [42] M. Omidi, H.R. Shamlouei, M. Noormohammadbeigi, The influence of Sc doping on structural, electronic and optical properties of Be₁₂O₁₂, Mg₁₂O₁₂ and Ca₁₂O₁₂ nanocages: a DFT study, J. Mol. Model. 23 (2017) 82.
- [43] H. Jouypazadeh, H. Farrokhpour, DFT and TD-DFT study of the adsorption and detection of sulfur mustard chemical warfare agent by the C₂₄, C₁₂Si₁₂, Al₁₂N₁₂, Al₁₂P₁₂, Be₁₂O₁₂, B₁₂N₁₂ and Mg₁₂O₁₂ nanocages, J. Mol. Struct. 1164 (2018) 227–238.
- [44] M. Solimannejad, S. Kamalinahad, E. Shakerzadeh, Silicon carbide nanotubes (SiCNTs) serving for catalytic decomposition of toxic diazomethane (DAZM) gas: a DFT study, Mol. Phys. 116 (2018) 414–422.
- [45] W. Kohn, L.J. Sham, Self-consistent equations including exchange and correlation effects, Phys. Rev. 140 (1965) A1133.
- [46] P. Hohenberg, W. Kohn, Inhomogeneous electron gas, Phys. Rev. 136 (1964) B864.
 [47] T. Yanai, D.P. Tew, N.C. Handy, A new hybrid exchange-correlation functional using the Coulomb-attenuating method (CAM-B3LYP), Chem. Phys. Lett. 393 (2004) 51–57.
- [48] Y. Zhao, D.G. Truhlar, The M06 suite of density functionals for main group thermochemistry, thermochemical kinetics, noncovalent interactions, excited states, and transition elements: two new functionals and systematic testing of four M06-class functionals and 12 other functionals, Theor. Chem. Acc. 120 (2008) 215–241.
- [49] J.-D. Chai, M. Head-Gordon, Long-range corrected hybrid density functionals with damped atom-atom dispersion corrections, PCCP 10 (2008) 6615–6620.
- [50] S. Grimme, J. Antony, S. Ehrlich, H. Krieg, A consistent and accurate ab initio parametrization of density functional dispersion correction (DFT-D) for the 94 elements H-Pu, J. Chem. Phys. 132 (2010), 154104.

M.D. Mohammadi et al.

- [51] T.V. Russo, R.L. Martin, P.J. Hay, Density functional calculations on first-row transition metals, J. Chem. Phys. 101 (1994) 7729–7737.
- [52] R. Dennington, T.A. Keith, J.M. Millam, GaussView, version 6.0. 16, Semichem Inc. Shawnee Mission KS, 2016.
- [53] J. Baker, An algorithm for geometry optimization without analytical gradients, J. Comput. Chem. 8 (1987) 563–574.
- [54] M.J. Frisch, G.W. Trucks, H.B. Schlegel, G.E. Scuseria, M.A. Robb, J.R. Cheeseman, G. Scalmani, V. Barone, G.A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A.V. Marenich, J. Bloino, B.G. Janesko, R. Gomperts, B. Mennucci, H.P. Hratchian, J.V. Ortiz, A.F. Izmaylov, J.L. Sonnenberg, Williams, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V.G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J.A. Montgomery Jr., J.E. Peralta, F. Ogliaro, M.J. Bearpark, J.J. Heyd, E.N. Brothers, K.N. Kudin, V.N. Staroverov, T.A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A.P. Rendell, J.C. Burant, S.S. Iyengar, J. Tomasi, M. Cossi, J.M. Millam, M. Klene, C. Adamo, R. Cammi, J.W. Ochterski, R.L. Martin, K. Morokuma, O. Farkas, J.B. Foresman, D.J. Fox, Gaussian 16 Rev. C.01, Wallingford, CT, 2016.
- [55] T. Lu, F. Chen, Multiwfn: A multifunctional wavefunction analyzer, J. Comput. Chem. 33 (2012) 580–592.
- [56] N.M. O'boyle, A.L. Tenderholt, K.M. Langner, Cclib: A library for packageindependent computational chemistry algorithms, J. Comput. Chem. 29 (2008) 839–845.
- [57] S.F. Boys, F. Bernardi, The calculation of small molecular interactions by the differences of separate total energies. Some procedures with reduced errors, Mol. Phys. 19 (1970) 553–566.

- Computational and Theoretical Chemistry 1198 (2021) 113168
- [58] J. Foresman, A. Frisch, Exploring Chemistry With Electronic Structure Methods: A Guide to Using Gaussian, Gaussian, Inc, Pittsburgh, PA, 1996.
- [59] J.-L. Bredas, Mind the gap!, Mater. Horiz. 1 (2014) 17–19.
- [60] F. Weinhold, Discovering chemistry with natural bond orbitals, John Wiley & Sons, 2012.
- [61] R.S. Mulliken, Electronic population analysis on LCAO-MO molecular wave functions. I, J. Chem. Phys. 23 (1955) 1833–1840.
- [62] I. Mayer, Improved definition of bond orders for correlated wave functions, Chem. Phys. Lett. 544 (2012) 83–86.
- [63] O.V. Sizova, L.V. Skripnikov, A.Y. Sokolov, Symmetry decomposition of quantum chemical bond orders, J. Mol. Struct. (Thoechem) 870 (2008) 1–9.
- [64] I. Mayer, P. Salvador, Overlap populations, bond orders and valences for 'fuzzy'atoms, Chem. Phys. Lett. 383 (2004) 368–375.
- [65] T. Lu, F. Chen, Bond order analysis based on the Laplacian of electron density in fuzzy overlap space, J. Phys. Chem. A 117 (2013) 3100–3108.
- [66] C.F. Matta, Hydrogen-Hydrogen Bonding: The Non-Electrostatic Limit of Closed-Shell Interaction Between Two Hydro, Hydrogen Bonding-New Insights, Springer, 2006, pp. 337–375.
- [67] H.J. Bohórquez, R.J. Boyd, C.F. Matta, Molecular model with quantum mechanical bonding information, J. Phys. Chem. A 115 (2011) 12991–12997.
- [68] E.R. Johnson, S. Keinan, P. Mori-Sánchez, J. Contreras-García, A.J. Cohen, W. Yang, Revealing noncovalent interactions, J. Am. Chem. Soc. 132 (2010) 6498–6506.
- [69] J. Contreras-García, E.R. Johnson, S. Keinan, R. Chaudret, J.-P. Piquemal, D. N. Beratan, W. Yang, NCIPLOT: a program for plotting noncovalent interaction regions, J. Chem. Theory Comput. 7 (2011) 625–632.